## EFFECT OF PEIERLS TRANSITION ON THE PHONON SPECTRUM IN ORGANIC CRYSTALS OF TTT<sub>2</sub>I<sub>3</sub> FOR DIFFERENT VALUES OF CARRIER CONCENTRATION

## Silvia ANDRONIC, Anatolie CASIAN, Ionel SANDULEAC, Ion BALMUS

Technical University of Moldova, Stefan cel Mare Avenue 168, Chisinau, Republic of Moldova

\*Corresponding author: Silvia Andronic, silvia.andronic@mt.utm.md

The metal-insulator transition of Peierls type and its effect on the phonon spectrum in quasi-one-dimensional organic crystals of  $TTT_2I_3$  for different values of carrier concentration is studied in 2D approximation. In the frame of the physical model, two electron-phonon interaction mechanism are taken into account. The first mechanism is of deformation potential type and the second one is of polaron type. The ratio of amplitudes of polaron-type interaction to the deformation potential one in *x* and *y* directions is characterized by parameters  $\gamma_1$  and  $\gamma_2$ . The interaction of carriers with structural defects is also considered. It was demonstrated that this interaction is very important to explain the behavior of Peierls transition. The method of retarded temperature dependent Green function is applied.

Computer modellings for 2D physical model of the crystal were performed for different values of carrier concentration. The numerical calculations for renormalized phonon spectrum,  $\Omega(q_x)$  as function of  $q_x$ , for different temperatures are presented in two cases: 1) when the interaction between transversal chains is neglected ( $q_y = 0$ ) and 2) when the interaction between the adjacent chains is considered ( $q_y \neq 0$ ). It is shown that in the second case when the interaction between adjacent chains is considered, the Peierls transition occurs at a lower temperature compared to the case when the interaction between transversal chains is neglected, for the same value of carrier concentration. Furthermore, it is shown that the Peierls structural transition strongly depends on iodine concentration. It was observed that with an increase in carrier concentration the Peierls critical temperature decreases.

**Keywords:** *Peierls transition, quasi-one-dimensional organic crystals, renormalized phonons, Peierls critical temperature, interchain interaction.* 

## References

- 1. ANDRONIC, S., CASIAN, A. Metal-Insulator Transition of Peierls Type in Quasi-One-Dimensional Crystals of TTT<sub>2</sub>I<sub>3</sub>. In: *Advances in Materials Physics and Chemistry*, 2017, 7, pp 212-222. https://doi.org/10.4236/ampc.2017.75017
- 2. ANDRONIC, S., BALMUS, I., CASIAN, A. Metal-Insulator Transition of Peierls Type in TTT<sub>2</sub>I<sub>3</sub> Crystals in 3D Approximation. *ICMCS 2017*, pp. 539-542.
- 3. ANDRONIC, S., SANDULEAC, I., CASIAN, A. Peierls Structural Transition in Organic Crystals of TTT<sub>2</sub>I<sub>3</sub> with Intermediate Carrier Concentration. In: *Springer Nature Switzerland AG 2020, I. Tiginyanu et al. (eds.), 4<sup>th</sup> International Conference on Nanotechnologies and Biomedical Engineering, IFMBE Proceedings, 2019, 77, pp. 199-202. https://doi.org/10.1007/978-3-030-31866-6\_40*